

Image® 70 DG Herbicide



Summer Weeds Controlled By Image.

Image controls a variety of summer weeds, including:

- >> Field Sandbur
- >> Perennial/Annual Ryegrass - overseeded
- >> Globe Sedge
- >> Green Kyllinga
- >> Rice Flatsedge
- >> Purple Nutsedge
- >> Yellow Nutsedge

Image also aids in controlling and reducing competition from:

- >> Dollarweed
- >> Tall Fescue
- >> Violets
- >> Annual Sedge
- >> Virginia Buttonweed

Winter Weeds Controlled By Image.

In addition to many summer weeds, **Image** also delivers effective control of many troublesome winter weeds. Applications should be timed to follow the first frost. With the exceptions of wild garlic and wild onion, applications can also be made immediately prior to or following the emergence of the following:

- >> Black Medic
- >> Buttercup
- >> Common Chickweed
- >> Mouseear Chickweed
- >> Cutleaf Evening Primrose
- >> Carolina Geranium
- >> Dovetail Geranium
- >> Hairy Bittercress
- >> Henbit
- >> Knawel
- >> Lawn Burweed
- >> Parsley Piert
- >> Purple Deadnettle
- >> Red Sorrel
- >> White Cover
- >> Wild Garlic
- >> Wild Onion

How To Apply Image.

Image should be applied at a rate of 0.2 ounces to 0.26 ounces per 1000 square feet (8.6 ounces to 11.4 ounces per acre). The high-use rate should be practiced where heavy weed infestation occurs.

For spot spraying with a handheld or backpack sprayer, mix 0.5 ounces of **Image** per 3 gallons of water and spray to wet. Approximately 1.5 gallons of spray solution will treat 1000 sq. ft.

Uniformly apply **Image** and avoid overlaps when spraying. Avoid drift onto vegetables, flowers, ornamental shrubs, and other desirable plants or injury may result. Irrigation or rainfall of 0.25 to 0.5 inches is suggested within 1 to 7 days of application to wash **Image** into the shoot/root zone where the greatest weed control activity occurs.

Image can also be applied without harm around and over a wide variety of ornamentals. For a complete list, consult the product label.